

## THE EFFECT OF DIFFERENT COMPOST: SOIL RATES ON SOME PARAMETERS OF THE SOIL-PLANT SYSTEM

Anita SZABO\*, Emese Berta SZABO, Marianna SIPOS, Imre VAGO

University of Debrecen, Debrecen, Hungary

Faculty of Agricultural and Food Sciences and Environmental Management,  
Department of Agricultural Chemistry and Soil Science

**ABSTRACT.** Composting is an alternative way for practicing site-specific and environmental friendly plant nutrient supply. Our aim was to study the effect of different composts on plant and soil system. Pot experiment with acidic sandy soil blended with three composts in five (0%, 5%, 10%, 25% and 50%) proportion was set up. Our experimental plant was perennial ryegrass (*Lolium perenne* L.). After the harvest of ryegrass we measured the fresh and dry weight of harvested plants, the P-, K-, Mg-contents, and the pH of soils. The three composts had different effect on dry weight production and on nutrient-supply of soil. We established that one of the compost had significantly larger effect on the dry weight of ryegrass compared to the others. In this study we proved that favorable compost/soil proportion is different in the case of different composts.

**Keywords:** sustainable plant production, compost mixing ratio

### INTRODUCTION

The location of waste emitted in increasing volume is one of the most important environmental problems of sustainable development. The disposal of by-products and waste materials formed in the public spaces of cities and during the processing of agricultural row material in a landfill is inconceivable, so we have to recycle them (Simándi, 2008). One of the solutions for this problem is composting. Beyond this the fertilization of horticultural crops with the composted waste materials, containing a large amount of organic matter content, is one of the alternative, environmental friendly plant nutrient supply practices. Scientific research on the positive effect of compost on soil parameters and yield has been published widely both in Hungarian and international literature (Gigliotti et al., 1966, Kádár-Morvai, 2007, Keserű, 2007). The reason of that we chose the more accurate cognition of compost utilization is to do more effective the site-specific nutrient supply. Our aim was to study and

evaluate the relations between the dry matter productions of plant, the compost:soil ration and the nutrient content of soil.

### MATERIALS AND METHODS

Compost utilization experiment with 3 different composts under controlled conditions was set up in the glasshouse of the Department of Agricultural Chemistry and Soil Science. Our aim was to study the effect of different composts on plant and soil system. We received the 3 composts from one of the partners of University of Debrecen in 2009. The composition and the production method of them are confidential. Composts were sieved (< 2 mm), because degradation of the large particles in the pots is slow. Composts were mixed with acidic sandy soil in four proportions (5%, 10%, 25% and 50%), in four repetition (*Table 1*). After the volumetric mixture we set up the pots randomized.

Table 1

The compost-soil ratio of treatments

Treatments	Compost (%)	Sandy soil (%)
1.	0	100
2.	5	95
3.	10	90
4.	25	75
5.	50	50

After one week maturation of compost-soil mixture we sowed perennial ryegrass (*Lolium perenne* L.). The advantage of ryegrass is that it grows quite fast, tolerates the glasshouse conditions well and it indicates the effect of treatments well. After the shooting of ryegrass the water supply of the 2.5 kg pots was carried

out at 60 per cent of field water capacity of soil. In our department previous researches confirmed that irrigation at 60 per cent of field water capacity is optimal in sandy soils (Loch et al., 1992). After the harvest we measured the fresh- and dry mass of ryegrass. We investigated the 0.01 M dm<sup>-3</sup> CaCl<sub>2</sub>

\*Correspondence: Anita Szabó, University of Debrecen, Faculty of Agricultural and Food Sciences and Environmental Management, Department of Agricultural Chemistry and Soil Science, no. 138. Böszörményi St., 4032 Debrecen, Hungary,  
Email: szaboanita@agr.unideb.hu  
Article received: May 2010; published: November 2010

extractable P-, K-, Mg-content and the pH values of dried and sieved soils.

### Chemical analysis

#### a.) 0.01 M CaCl<sub>2</sub>-soil extraction method

The P-, K-, Mg-contents of the soils were measured in 0.01 mol CaCl<sub>2</sub> extractant (Houba et al., 1990) with UNICAM SP95B AAS and SKALAR Continuous Flow Analyzer at the laboratory of the Department of Agricultural Chemistry and Soil Science. We measured the pH of the soils after 0.01 M CaCl<sub>2</sub> extraction.

#### b.) Statistical analyses

We used variance analyses for the processing of the data. All statistical analyses were performed with a Microsoft Excel Macro (Tolner, 2008, Vágó, 2008) according to Sváb (1981). With this program we determined the significance level of the treatment effect and significant difference in P = 5 % (SD5%).

## RESULTS AND DISCUSSION

The fresh- and dry weights of ryegrass in each treatments (5 compost: soil rate; 3 compost) are shown in Figure 1 and Figure 2.




Fig. 1 Fresh weight production of ryegrass in each treatment (g 2.5 kg<sup>-1</sup>)

As it is represented in Figure 1 and Figure 2, the compost/soil ratio and the different composts had a significant effect (P = 0.1 %) on the fresh and dry weight of ryegrass.

We found the highest positive effect in the case of 1. compost. If a 25 % compost dose were applied, the fresh weight increasing effect of 1. compost would reach a value of 45 g pot<sup>-1</sup>. The increasing dose of compost didn't caused higher yields. It can be stated that the optimal compost:soil ratio is 25:75 %. The 25% and 50% dose of 3. compost increased significantly the yield. The yield increasing effect of 50 % compost was nearly 10 g pot<sup>-1</sup>. The 2. compost had a moderate effect, but the 10% compost dose increased significantly the dry and fresh weight of ryegrass. As the data show the recommended compost dose is different for each compost.

The P-, K-, Mg content of the soil determined in 0.01 M CaCl<sub>2</sub> are represented in Figure 3., 4., 5. The compost:soil ratio modified significantly (P = 0.1 %) the CaCl<sub>2</sub> extractable nutrient content.

All studied compost increased significantly the amount of CaCl<sub>2</sub>-P (Figure 3.). The applying of 50 % compost increased the P content of soil by 2-4 times higher. We found the highest easily available P concentration (11 mg kg<sup>-1</sup>) in case of 1. compost. A moderate effect was observable with the 2. and 3. compost.

As it is represented in Figure 4. and Figure 5. the increasing compost rates caused the increasing of K, Mg content of soil. The CaCl<sub>2</sub>-K and -Mg content of the 2. compost was higher than that of other composts. We found the lowest nutrient content in case of 3. compost.

The pH values of the soils determined in CaCl<sub>2</sub> are shown in Figure 6. The treatments increased significantly the pH values of soils (P = 0.1 %). The 2. compost had the highest effect on the pH. All ratios caused significant increase. If a 50 % compost dose were applied the pH would be in neutral range (pH = 6.81). In case of 1. and 3. composts the 25 and 50 % dose caused significant increase.




Fig. 2 Dry weight production of ryegrass in each treatment (g 2.5 kg<sup>-1</sup>)




Fig. 3 The 0.01 M dm<sup>-3</sup> CaCl<sub>2</sub> extractable P content of soil




Fig. 4 The 0.01 M dm<sup>-3</sup> CaCl<sub>2</sub> extractable K content of soil




Fig. 5 The 0.01 M dm<sup>-3</sup> CaCl<sub>2</sub> extractable Mg content of soil




Fig. 6 pH(CaCl<sub>2</sub>) of soil

## CONCLUSIONS

From the results of the study it was concluded as follows: However the nutrient content and pH of soils increased significantly as an effect of increasing compost doses, the optimal compost dose in respect of fresh and dry weight production was different in case of each compost. The 1. compost affected the highest increase in the amount of yield, and the optimal compost:soil rate was 25:75 %. The optimal dose of 3. compost was the 50 % rate. The 2. compost had a moderate effect, but the 10 % compost dose increased significantly the dry and fresh weight of ryegrass.

## REFERENCES

- Gigliotti, G., Businelli, D. & Giusquiani, P.L. (1966): Trace metal uptake and distribution in corn plants grown on a 6-year urban waste compost amended soil. *Agric. Ecosyst. and Environm.* 58. 199–206.
- Houba, V.J.G. – Novozamsky, L. – Lexmond, T.M. – Van der Lee, J.J. (1990): Applicability of 0,01 M CaCl<sub>2</sub> as a single extraction solution for the assessment of the nutrient status of soils and other diagnostic purposes. *Commun. Soil Sci. Plant Anal.* 21. 2281-2290. p.
- Kádár, I. – Morvai, B. (2007): Ipari-kommunális szennyvíziszap-terhelés hatásának vizsgálata tenyészedény-kísérletben. *AGROKÉMIA ÉS TALAJTAN* 56 (2007) 2
- Keserű, Zs. (2007): A szennyvíziszap-komposzt erdészeti hasznosíthatóságának kérdései.

Erdészeti, Környezettudományi, Természetvédelmi és Vadgazdálkodási Tudományos Konferencia (EKTV-TK) konferencia 2007. december 11. Sopron.

- Loch, J. - Kiss Sz. - Vágó I. (1992): A kálium-, kalcium-, magnézium- és vízellátás hatása az őszi búza szemtermésére és magnéziumfelvételére. 4. Magyar Magnézium Szimpózium, Balatonszéplak. In: *Magnesium Research* 5. Abstr. 238.
- Simándi, P. (2008): Különböző szerves hulladékok és kezelésük után keletkezett termékek kémiai vizsgálata. *Értekezés, Debrecen.* 8 p.
- Sváb, J. (1981): *Biometriai módszerek a kutatásban.* Mezőgazdasági Kiadó, Budapest
- Tolner, L. – Vágó, I. – Czinkota, I. – Rékási, M. – Kovács, Z. (2008): Field testing of a new, more efficient liming method. *Cereal Research Communications.* 36. 543 - 546. p.
- Vágó, I. – Tolner, L. – Eichler-Löbermann, B. – Czinkota, I. – Kovács, B. (2008): Long-term effects of liming on the dry matter production and chemical composition of perennial ryegrass (*Lolium perenne* L.). *Cereal Research Communications.* 36. 103 - 106. p.